
Scleria robusta, a New Name for *Scleria grandis* Core (Cyperaceae)

Konraed Camelbeke and Paul Goetghebeur

Gent University, Department of Biology, Laboratory of Botany, K.L. Ledeganckstraat 35,
B-9000 Gent, Belgium

ABSTRACT. *Scleria robusta* Camelbeke & Goetghebeur is proposed as a new name for the later homonym *Scleria grandis* Core.

When Core (1945) published a new neotropical species of *Scleria* (Cyperaceae), naming it *Scleria grandis* Core, he overlooked the fact that this combination had already been used by Labillardière (1800) for the Australian species now generally recognized as *Gahnia grandis* (Labillardière) S. T. Blake (see, e.g., Wilson, 1994: 281). It was indeed Blake (1969: 33) who made the correct combination in *Gahnia*, including the homotypic synonym *Gahnia psittacorum* Labillardière. However, he did not propose a new name for *Scleria grandis* Core, and therefore an avowed substitute for this later homonym must be proposed.

Scleria robusta Camelbeke & Goetghebeur, nom. nov. Replaced name: *Scleria grandis* Core, J. Wash. Acad. Sci. 35(10): 322. 1945, not Labillardière, Voy. Rech. Pérouse: 146. 1800. TYPE: Colombia. Vaupés: Selva del Tui-Igarape, cerca de Mitú, 200 m, 17 Sep. 1939, J. Cuatrecasas 6870 (lectotype, selected here, US 1795819; isolectotype, US 1795820).

This distinctive member of section *Hymenolymrum* is distinguished from its nearest relative, *Scleria cyperina* Kunth, by the broader leaves (18–

45 vs. 7–15 mm), the broader foliaceous primary bracts (18–35 vs. 6–13 mm), the smooth nutlet, and the more robust habit of the former; hence *robusta* as a new epithet.

Paratypes. COLOMBIA. Caquetá: Cerro de La Sardina, Florencia, bosque, ca. 500 m, 30 Mar. 1940, J. Cuatrecasas 8890 (US). BRAZIL. Amazonas: Santa Isabel, Rio Negro, 15–16 Feb. 1944, J. T. Jr. Baldwin 3421 (US 2 sheets).

Acknowledgments. We thank Paul E. Berry (WIS) for bringing this nomenclatural problem to our attention. Karen L. Wilson (NSW) is thanked for the fruitful discussion and help. The US curator is acknowledged for loan of the type specimens of *Scleria grandis* Core, and the FI curator for providing photos of the Labillardière material. The librarians of BR and NY are thanked for providing photocopies of the requisite old literature. Tony Reznicek (MICH) and an anonymous reviewer are thanked for useful comments on the manuscript.

Literature Cited

- Blake, S. T. 1969. Studies in Cyperaceae. Contr. Queensland Herb. 8: 1–48.
Core, E. L. 1945. Two new species of *Scleria* from the Upper Amazon Valley. J. Wash. Acad. Sci. 35: 322.
Labillardière, J. J. H. de. 1800. Relation du Voyage à la Recherche de la Pérouse, Vol. 1. Paris.
Wilson, K. L. 1994. Cyperaceae. Pp. 238–356 in N. G. Walsh & T. J. Entwistle (editors), Flora of Victoria, Vol. 2. Inkata Press, Melbourne.